


Amendments to the Claims

This Listing of Claims will replace all prior versions, and listings, of claims in the specification:

Listing of Claims:

1.-6. (Canceled)

7. (Currently Amended) ~~The A~~ compound according to claim 3 of having the formula



wherein

R₁ and R₂ are independently hydrogen, cyan, halo, nitro, optionally substituted amino, lower alkyl C₁₋₄ alkyl, trifluoromethyl, -CO₂H, CO₂C₁₋₄ alkyl, C(O)NHC₁₋₄ alkyl, or lower C₁₋₄ alkoxy, or

R₁ and R₂ combined together with the carbon atoms to which they are attached form an optionally substituted 6-membered aromatic ring;

W is -NR₅C(O)R₆, -NR₅C(O)OR₆, -NR₅C(O)NR₆R₇, -NR₅C(S)NR₆R₇, -NR₅S(O)₂R₆, -NR₅R₈, -C(O)NR₆R₇, -OR₉ or -OC(O)NR₆R₇ in which

R₅ and R₇ are independently hydrogen or methyl; or

R₅ and R₁ are alkylene which combined together with the nitrogen atom to which R₅ is attached and the carbon atoms to which W and R₁ are attached form a 5-membered ring;

R₆ is optionally substituted alkyl, aryl, heteroaryl, cycloalkyl, aralkyl or heteroaralkyl;

R₈ is optionally substituted alkyl, aralkyl or heteroaralkyl;

R₉ is hydrogen, optionally substituted alkyl, aralkyl, heteroaralkyl or alkanoyl; or

W is aryl or heteroaryl; or

W is hydrogen provided that R₁ is NR₅Z in which Z is -C(O)R₆, -C(O)OR₆, -C(O)NR₆R₇, -C(S)NR₆R₇, -S(O)₂R₆, or R₈; or

W and R₁ combined together with the carbon atoms they are attached to form a 6-membered aromatic ring optionally substituted with alkyl, alkoxy, aryl, heteroaryl, halo, -NR₅Z, -C(O)NR₆R₇, -OR₉ or -OC(O)NR₆R₇;

X is CH;

Y is CH or nitrogen; or

~~X=Y is CH₂, oxygen, sulfur or NR₁₀ in which R₁₀ is hydrogen or methyl;~~

R₁₃ and R₁₄ are independently hydrogen, hydroxy or optionally substituted lower C₁₋₄ alkyl; or

a pharmaceutically acceptable salt thereof.

8. (Currently Amended) The compound according to claim 7 wherein

R₁ is hydrogen;

R₂ is hydrogen, chloro, methoxy, ethoxy, propoxy or optionally substituted amino;

W is -NR₅C(O)R₈, -NR₅C(O)OR₆, -NR₅C(O)NR₆R₇, -NR₅C(S)NR₆R₇, -NR₅S(O)₂R₆, -NR₅R, -C(O)NR₆R₇, -OR₉ or -OC(O)NR₆R₇ in which

R₅ and R₇ are independently hydrogen or methyl;

R₆ is optionally substituted alkyl, aryl, heteroaryl, cycloalkyl, aralkyl or heteroaralkyl;

R₈ is optionally substituted alkyl, aralkyl or heteroaralkyl;

R₉ is hydrogen, optionally substituted alkyl, aralkyl, heteroaralkyl or alkanoyl;

X is CH;

Y is CH;

R₁₃ and R₁₄ are independently hydrogen, hydroxy or optionally substituted lower alkyl; or a pharmaceutically acceptable salt thereof.

9. (Currently Amended) The compound according to claim 7 wherein

R₁ is methyl, methoxy or optionally substituted amino;

R₂ is hydrogen;

W is -NR₅C(O)R₆, -NR₅C(O)OR₆, -NR₅C(O)NR₆R₇, -NR₅C(S)NR₆R₇, -NR₅S(O)₂R₆, -NR₅R₈, -C(O)NR₆R₇, -OR₉ or -OC(O)NR₆R₇ in which

R_5 and R_7 are independently hydrogen or methyl;

R_6 is optionally substituted alkyl, aryl, heteroaryl, cycloalkyl, aralkyl or heteroaralkyl;

R_8 is optionally substituted alkyl, aralkyl or heteroaralkyl;

R_9 is hydrogen, optionally substituted alkyl, aralkyl, heteroaralkyl or alkanoyl;

X is CH;


Y is CH;

R_{13} and R_{14} are independently hydrogen, hydroxy or optionally substituted lower alkyl; or a pharmaceutically acceptable salt thereof.

10. (Canceled)

11. (Canceled)

12. (Currently Amended) The compound according to claim 7 of the formula



wherein

W is $-NR_5C(O)R_6$, $-NR_5C(O)OR_6$, $-NR_5C(O)NR_6R_7$, $-NR_5C(S)NR_6R_7$, $-NR_5S(O)_2R_6$, $-NR_5R_8$, $-C(O)NR_6R_7$, $-OR_9$ or $-OC(O)NR_6R_7$ in which

R_5 and R_7 are independently hydrogen or methyl;

R_6 is optionally substituted alkyl, aryl, heteroaryl, cycloalkyl, aralkyl or heteroaralkyl;

R_8 is optionally substituted alkyl, aralkyl or heteroaralkyl;


R_9 is hydrogen, optionally substituted alkyl, aralkyl, heteroaralkyl or alkanoyl;

Y is CH;

R_{13} and R_{14} are independently hydrogen, hydroxy or optionally substituted lower alkyl;

or a pharmaceutically acceptable salt thereof.

13. (Currently Amended) The A compound according to claim 7 of the formula



wherein

R₂ is hydrogen, halo or alkoxy;

Y is CH or nitrogen;

R₁₃ and R₁₄ are independently hydrogen, hydroxy or optionally substituted lower alkyl;

R₁₅ is hydrogen, -NR₅C(O)R₆, -NR₅C(O)OR₆, -NR₅C(O)NR₆R₇, -NR₅C(S)NR₆R₇, -NR₅S(O)₂R₆, -NR₅R₈, -C(O)NR₆R₇, -OR₉ or -OC(O)NR₆R₇ in which

R₅ and R₇ are independently hydrogen or methyl;

R₆ is optionally substituted alkyl, aryl, heteroaryl, cycloalkyl, aralkyl or heteroaralkyl;


R₈ is optionally substituted alkyl, aralkyl or heteroaralkyl;

R₉ is hydrogen, optionally substituted alkyl, aralkyl, heteroaralkyl or alkanoyl;

or a pharmaceutically acceptable salt thereof.

14-17. (Canceled).

18. (Currently Amended) The compound according to claim 3 of the formula



wherein

R₁ and R₂ are independently hydrogen, halo, optionally substituted amino, lower C₁₋₄ alkyl or lower C₁₋₄ alkoxy; or

R_1 and R_2 combined together form an optionally substituted 6-membered aromatic ring;
 W is $-NR_5C(O)R_6$, $NR_5C(O)OR_6$, $-NR_5C(O)NR_6R_7$, $-NR_5C(S)NR_6R_7$, $-NR_5S(O)_2R_6$,
 $-NR_5R_8$, $-C(O)NR_6R_7$, $-OR_9$ or $-OC(O)NR_6R_7$ in which

R_5 and R_7 are independently hydrogen or methyl; or

R_5 and R_1 are alkylene which combined together with the nitrogen atom to which
 R_5 is attached and the carbon atoms to which W and R_1 are attached form a 5-membered ring;

R_6 is optionally substituted alkyl, aryl, hetroaryl, cycloalkyl, aralkyl or
heteroaralkyl;

R_8 is optionally substituted alkyl, aralkyl or heteroaralkyl;

R_9 is hydrogen, optionally substituted alkyl, aralkyl, heteroaralkyl or alkanoyl; or

W is aryl or heteroaryl; or

W and R_1 combined together with the carbon atoms to which they are attached
form a 6-membered aromatic ring optionally substituted with alkyl, alkoxy, aryl, heteroaryl, halo,
 $-NR_5Z$, $-C(O)NR_6R_7$, $-OR_9$ or $-OC(O)NR_6R_7$ in which

Z is $-C(O)R_6$, $-C(O)OR_6$, $-C(O)NR_6R_7$, $-C(S)NR_6R_7$, $-S(O)_2R_6$, or $-R_8$;

R_{13} and R_{14} are independently hydrogen, hydroxy or optionally substituted lower alkyl;

X is CH;

Y is CH or nitrogen; or

~~$X=Y$ is CH_2 , oxygen, sulfur or NR_{10} in which R_{10} is hydrogen or methyl;~~

or a pharmaceutically acceptable salt thereof.

19. (Currently Amended) The compound according to claim 18 wherein

R_1 is hydrogen;

R_2 is hydrogen, chloro, methoxy, ethoxy, propoxy or optionally substituted amino;

W is $-NR_5C(O)R_6$, $-NR_5C(O)OR_6$, $-NR_5C(O)NR_6R_7$, $-NR_5C(S)NR_6R_7$, $-NR_5S(O)_2R_6$,
 $-NR_5R_8$, $-C(O)NR_6R_7$, $-OR_9$ or $-OC(C)NR_6R_7$ in which

R_5 and R_7 are independently hydrogen or methyl;

R_6 is optionally substituted alkyl, aryl, hetroaryl, cycloalkyl, aralkyl or
heteroaralkyl;

R_8 is optionally substituted alkyl, aralkyl or heteroaralkyl;


R_9 is hydrogen, optionally substituted alkyl, aralkyl, heteroaralkyl or alkanoyl;
X is CH;
Y is CH;
 R_{13} and R_{14} are independently hydrogen, hydroxy or optionally substituted lower alkyl;
or a pharmaceutically acceptable salt thereof.

20. (Currently Amended) The compound according to claim 18 wherein

R_1 is methyl, methoxy or optionally substituted amino;
 R_2 is hydrogen;
W is $-NR_5C(O)R_6$, $-NR_5C(O)OR_6$, $-NR_5C(O)NR_6R_7$, $-NR_5C(S)NR_6R_7$, $-NR_5S(O)_2R_6$,
 $-NR_5R_8$, $-C(O)NR_6R_7$, $-OR_9$ or $-OC(O)NR_6R_7$, in which
 R_5 and R_7 are independently hydrogen or methyl;
 R_8 is optionally substituted alkyl, aryl, heteroaryl, cycloalkyl, aralkyl or
heteroaralkyl;
 R_9 is optionally substituted alkyl, aralkyl or heteroaralkyl;
 R_9 is hydrogen, optionally substituted alkyl, aralkyl, heteroaralkyl or alkanoyl;
X is CH;
Y is CH;
 R_{13} and R_{14} are independently hydrogen, hydroxy or optionally substituted lower alkyl;
or a pharmaceutically acceptable salt thereof.

21. (canceled)

22. (Currently Amended) The compound according to claim 18 of the formula



W is -NR₅C(O)R₆, -NR₅C(O)OR₆, -NR₅C(O)NR₆R₇, -NR₅C(S)NR₆R₇, -NR₅S(O)₂R₆, -NR₅R₈, -C(O)NR₆R₇, -OR₉ or -OC(O)NR₆R₇ in which

R₅ and R₇ are independently hydrogen or methyl;

R₆ is optionally substituted alkyl, aryl, heteroaryl, cycloalkyl, aralkyl or heteroaralkyl;

R₈ is optionally substituted alkyl, aralkyl or heteroaralkyl;

R₉ is hydrogen, optionally substituted alkyl, aralkyl, heteroaralkyl or alkanoyl;

Y is CH;

R₁₃ and R₁₄ are independently hydrogen, hydroxy or optionally substituted lower alkyl; or a pharmaceutically acceptable salt thereof.

23-24. (Canceled)

25. (Withdrawn) A method for the inhibition of 11 β -hydroxysteroid dehydrogenase type 1 (11 β -HSD1) oxoreductase activity in mammals, which method comprises administering to a mammal in need thereof a therapeutically effective amount of a compound of claim 1.
26. (Withdrawn) A method to control glucocorticoid concentration in mammals which method comprises administering to a mammal in need thereof a therapeutically effective amount of a compound of claim 1.
27. (Withdrawn) A method according to claim 26, which comprises lowering intracellular and hepatic glucocorticoid concentrations, increasing insulin sensitivity in the adipose tissue and in the muscle, reducing lipolysis and free fatty acid production in the adipose tissue, and inhibiting hepatic gluconeogenesis.
28. (Withdrawn) A method for the treatment of conditions associated with 11 β -HSD1 oxoreductase activity in mammals which comprises administering to a mammal in need thereof a therapeutically effective amount of a compound of claim 1.
29. (Withdrawn) A method for the treatment of glucocorticoid associated disorders in mammals which method comprises administering to a mammal in need thereof a therapeutively effective amount of a compound of claim 1.

30. (Withdrawn) A method according to claim 29, which comprises administering a compound of claim 1 in combination with a therapeutically effective amount of insulin, insulin derivative or mimetic, insulin secretagogue, insulinotropic sulfonylurea receptor ligand, insulin sensitizer, biguanide, alpha-glucosidase inhibitor, GLP-1, GLP-1 analog or mimetic, DPP-IV inhibitor, hypolipidemic agent, anti-obesity agent, cholestyramine, fibrate, nicotinic acid, or aspirin.

31. (Withdrawn) A method for the treatment of impaired glucose tolerance in Type 2 diabetes which method comprises administering to a mammal in need thereof a therapeutically effective amount of a compound of claim 1.

32. (Withdrawn) A method for the treatment of Syndrome-X, dyslipidemia, hypertension and central obesity which method comprises administering to a mammal in need thereof a therapeutically effective amount of a compound of claim 1.

33. (Currently Amended) A pharmaceutical composition, comprising:

the compound of claim 1 preferably in a therapeutically effective amount, in combination with one or more pharmaceutically acceptable carriers.

34. (Canceled)

35. (Canceled)

36. (Canceled)

37. (Canceled)

38. (Canceled)

39. (Canceled)